

INDIVIDUAL ANTS SPECIALIZE ON PARTICULAR APHID HERDS  
(HYMENOPTERA: FORMICIDAE; HOMOPTERA: APHIDIDAE)

BERT C. EBBERS AND EDWARD M. BARROWS

(BCE) 2415 East G Avenue, Kalamazoo, Michigan 49004; (EMB) Department of Biology, Georgetown University, Washington, D.C. 20057.

---

*Abstract.*—Three ant-aphid associations were each observed for ten consecutive days. Eighty-one percent of 103 individually-marked ant foragers visited only one of many conspecific aphid groups on a particular plant. Specialization on particular herds may help ants maximize their net energy gain from aphid honeydew.

---

Ant species have coevolved to different degrees with trophobionts, homopterans and lepidopterans from which they acquire food (Wilson, 1971). For example, ant tending and obtaining "honeydew" from aphids is commonly seen in many habitats and has been studied numerous times (see Auclair, 1963, and Way, 1963, for authoritative reviews of this symbiosis). Because there appear to be no reports of individual ant foragers specializing on particular herds of honeydew-producing aphids, we relate our discoveries regarding this specialization and speculate on its adaptive significance.

Three associations of ants and aphids were observed from 29 June to 4 August 1978 on the property of the University of Michigan Biological Station, Cheboygan County, Michigan. Association 1 was the ant *Formica subnuda* Emery tending the aphids *Pterocomma* sp. on branches and *Chaitophorus nigrae* Oestlund on leaves of *Salix interior* Rowlee. Association 2 was *Formica subsericea* Say tending *C. nigrae* on leaves of *S. interior*. Association 3 was *Camponotus noveboracensis* (Fitch) tending *Chaitophorus* sp. on petioles of *Populus tremuloides* Michx. Each association was examined for ten consecutive days from 0215 to 2300 hours and from one to four times ( $\bar{x} = 2$ ) per day. Ants were removed from aphid herds and marked for individual recognition on their thoraces and gasters with fast-drying enamel paints and replaced on herds within 3 min.

A total of 158 ants was marked in the three associations; 65% of them (103) were seen again on at least one of the ten days during which associations were observed. Eighty-one percent of the 103 ants were seen at only one of all conspecific herds on particular plants. Ten percent changed herds once; 6%, twice; and 3%, thrice. The ants showed a mean herd constancy



( $P > 0.05$ ). Association 3 ants may have moved to closest herds based on distances over leaves rather than stems.

An experiment was attempted to examine the possibility that ants return to their original herds after being artificially removed from them and placed at new herds. However, the ants were extremely wary and jumped off plants when they were approached within 3 cm, or else they jumped off after they were placed on new herds. Their wariness may have resulted from their previous experiences with an investigator during marking.

In conclusion, this study reveals that individual ants of the same colony tend to visit only one or few of many herds of conspecific aphids on a particular plant for up to ten days and perhaps longer. This specialization on particular herds may be an integral part of such ants maximizing their net energy gains from aphids. By being constant to particular herds, these ants could move directly between such herds and their nests and avoid wasting time and energy in searching for untended aphids or competing with other aphid-tending ants. Further study is needed to elucidate the adaptive significance of this ant specialization; whether the ants recognize particular aphid herds or are constant to certain locations on aphid-harboring plants; why ants "decide" to tend one aphid herd rather than another; why ants change herds; and related subjects. The phenomenon of *Ortsreu*, patrolling only certain portions of a colonial feeding territory by a particular ant forager (Oster and Wilson, 1978), is likely to be related to aphid herd specialization. Finally, our investigation indicates that studies of ergonomic optimization in honeydew-collecting insects should take into consideration that foragers may specialize on particular groups of honeydew-producing insects.

#### ACKNOWLEDGMENTS

Manya B. Stoetzel (Systematic Entomology Laboratory USDA, Beltsville, Maryland) identified aphids; David R. Smith (Systematic Entomology Laboratory USDA, Washington, D.C.), ants. We are grateful for the opportunity to work at the University of Michigan Biological Station. Julia Johnson helped prepare the manuscript.

#### LITERATURE CITED

- Auclair, J. L. 1963. Aphid feeding and nutrition. *Ann. Rev. Entomol.* 8: 439-490.
- Oster, G. F. and E. O. Wilson. 1978. *Caste and ecology in the social insects*. Princeton Univ. Press, Princeton, N.J. 352 pp.
- Way, M. J. 1963. Mutualism between ants and honeydew-producing Homoptera. *Ann. Rev. Entomol.* 8: 307-344.
- Wilson, E. O. 1971. *The insect societies*. Belknap Press of Harvard Univ. Press, Cambridge, Mass. 548 pp.